Doctors Are The Third Leading Cause of Death in the US, Killing 225,000 People Every Year



- 12,000 -- unnecessary surgery
- 7,000 -- medication errors in hospitals
- 20,000 -- other errors in hospitals
- 80,000 -- infections in hospitals
- 106,000 -- non-error, negative effects of drugs

These total to 225,000 deaths per year from iatrogenic causes!!

What does the word iatrogenic mean? This term is defined as induced in a patient by a physician's activity, manner, or therapy. Used especially of a complication of treatment.

Dr. Starfield offers several warnings in interpreting these numbers:

- First, most of the data are derived from studies in hospitalized patients.
- Second, these estimates are for deaths only and do not include negative effects that are associated with disability or discomfort.
- Third, the estimates of death due to error are lower than those in the IOM report.

If the higher estimates are used, the deaths due to iatrogenic causes would range from 230,000 to 284,000. In any case, 225,000 deaths per year constitutes the third leading cause of death in the United States, after deaths from heart disease and cancer. Even if these figures are overestimated, there is a wide margin between these numbers of deaths and the next leading cause of death (cerebrovascular disease).

Another analysis concluded that between 4% and 18% of consecutive patients experience negative effects in outpatient settings, with:

- 116 million extra physician visits
- 77 million extra prescriptions
- 17 million emergency department visits
- 8 million hospitalizations
- 3 million long-term admissions
- 199,000 additional deaths
- \$77 billion in extra costs

The high cost of the health care system is considered to be a deficit, but seems to be tolerated under the assumption that better health results from more expensive care.



However, evidence from a few studies indicates that as many as 20% to 30% of patients receive inappropriate care.

An estimated 44,000 to 98,000 among them die each year as a result of medical errors.

This might be tolerated if it resulted in better health, but does it? Of 13 countries in a recent comparison, the United States ranks an average of 12th (second from the bottom) for 16 available health indicators. More specifically, the ranking of the US on several indicators was:

- 13th (last) for low-birth-weight percentages
- 13th for neonatal mortality and infant mortality overall
- 11th for postneonatal mortality
- 13th for years of potential life lost (excluding external causes)
- 11th for life expectancy at 1 year for females, 12th for males
- 10th for life expectancy at 15 years for females, 12th for males
- 10th for life expectancy at 40 years for females, 9th for males
- 7th for life expectancy at 65 years for females, 7th for males
- 3rd for life expectancy at 80 years for females, 3rd for males
- 10th for age-adjusted mortality

The poor performance of the US was recently confirmed by a World Health Organization study, which used different data and ranked the United States as 15th among 25 industrialized countries.

There is a perception that the American public "behaves badly" by smoking, drinking, and perpetrating violence." However the data does not support this assertion.

- The proportion of females who smoke ranges from 14% in Japan to 41% in Denmark; in the United States, it is 24% (fifth best). For males, the range is from 26% in Sweden to 61% in Japan; it is 28% in the United States (third best).
- The US ranks fifth best for alcoholic beverage consumption.
- The US has relatively low consumption of animal fats (fifth lowest in men aged 55-64 years in 20 industrialized countries) and the third lowest mean cholesterol concentrations among men aged 50 to 70 years among 13 industrialized countries.

These estimates of death due to error are lower than those in a recent Institutes of Medicine report, and if the higher estimates are used, the deaths due to iatrogenic causes would range from 230,000 to 284,000.

Even at the lower estimate of 225,000 deaths per year, this constitutes the third leading cause of death in the US, following heart

disease and cancer.

Lack of technology is certainly not a contributing factor to the US's low ranking.

- Among 29 countries, the United States is second only to Japan in the availability of magnetic resonance imaging units and computed tomography scanners per million population. 17
- Japan, however, ranks highest on health, whereas the US ranks among the lowest.
- It is possible that the high use of technology in Japan is limited to diagnostic technology not matched by high rates of treatment, whereas in the US, high use of diagnostic technology may be linked to more treatment.
- Supporting this possibility are data showing that the number of employees per bed (full-time equivalents) in the United States is highest among the countries ranked, whereas they are very low in Japan, far lower than can be accounted for by the common practice of having family members rather than hospital staff provide the amenities of hospital care.

Journal American Medical Association July 26, 2000;284(4):483-5

DR .MERCOLA'S



Dr. Mercola's Comments:

Folks, this is what they call a "Landmark Article". Only several ones like this are published every year. One of the major reasons it is so huge as that it is published in JAMA which is the largest and one of the most respected medical journals in the entire world.

I did find it most curious that the best wire service in the world, Reuter's, did not pick up this article. I have no idea why they let it slip by.

I would encourage you to bookmark this article and review it several times so you can use the statistics to counter the arguments of your friends and relatives who are so enthralled with the traditional medical paradigm. These statistics prove very clearly that the system is just not working. It is broken and is in desperate need of repair.

I was previously fond of saying that drugs are the fourth leading cause of death in this country. However, this article makes it quite clear that the more powerful number is that doctors are the third leading cause

of death in this country killing nearly a quarter million people a year. The only more common causes are cancer and heart disease.

This statistic is likely to be seriously underestimated as much of the coding only describes the cause of organ failure and does not address iatrogenic causes at all.

Japan seems to have benefited from recognizing that technology is wonderful, but just because you diagnose something with it, one should not be committed to undergoing treatment in the traditional paradigm. Their health statistics reflect this aspect of their philosophy as much of their treatment is not treatment at all, but loving care rendered in the home.

Care, not treatment, is the answer. Drugs, surgery and hospitals are rarely the answer to chronic health problems. Facilitating the God-given healing capacity that all of us have is the key. Improving the diet, exercise, and lifestyle are basic.

Effective interventions for the underlying emotional and spiritual wounding behind most chronic illness are also important clues to maximizing health and reducing disease.

Author/Article Information

Author Affiliation: Department of Health Policy and Management, Johns Hopkins School of Hygiene and Public Health, Baltimore, Md. Corresponding Author and Reprints: Barbara Starfield, MD, MPH, Department of Health Policy and Management, Johns Hopkins School of Hygiene and Public Health, 624 N Broadway, Room 452, Baltimore, MD 21205-1996 (e-mail: bstarfie@jhsph.edu).

References

1. Schuster M, McGlynn E, Brook R. How good is the quality of health care in the United States? Milbank Q. 1998;76:517-563.

2. Kohn L, ed, Corrigan J, ed, Donaldson M, ed. To Err Is Human: Building a Safer Health System. Washington, DC: National Academy Press; 1999.

3. Starfield B. Primary Care: Balancing Health Needs, Services, and Technology. New York, NY: Oxford University Press; 1998.

4. World Health Report 2000. Accessed June 28, 2000.

5. Kunst A. Cross-national Comparisons of Socioeconomic Differences in Mortality. Rotterdam, the Netherlands: Erasmus University; 1997.

6. Law M, Wald N. Why heart disease mortality is low in France: the time lag explanation. BMJ. 1999;313:1471-1480.

7. Starfield B. Evaluating the State Children's Health Insurance Program: critical considerations. Annu Rev Public Health. 2000;21:569-585.

8. Leape L.Unecessarsary surgery. Annu Rev Public Health. 1992;13:363-383.

9. Phillips D, Christenfeld N, Glynn L. Increase in US medication-error deaths between 1983 and 1993. Lancet. 1998;351:643-644.

10. Lazarou J, Pomeranz B, Corey P. Incidence of adverse drug reactions in hospitalized patients. JAMA. 1998;279:1200-1205.

11. Weingart SN, Wilson RM, Gibberd RW, Harrison B. Epidemiology and medical error. BMJ. 2000;320:774-777.

12. Wilkinson R. Unhealthy Societies: The Afflictions of Inequality. London, England: Routledge; 1996.

13. Evans R, Roos N. What is right about the Canadian health system? Milbank Q. 1999;77:393-399.

14. Guyer B, Hoyert D, Martin J, Ventura S, MacDorman M, Strobino D. Annual summary of vital statistics1998. Pediatrics. 1999;104:1229-1246.

15. Harrold LR, Field TS, Gurwitz JH. Knowledge, patterns of care, and outcomes of care for generalists and specialists. J Gen Intern Med. 1999;14:499-511.

16. Donahoe MT. Comparing generalist and specialty care: discrepancies, deficiencies, and excesses. Arch Intern Med. 1998;158:1596-1607.

17. Anderson G, Poullier J-P. Health Spending, Access, and Outcomes: Trends in Industrialized Countries. New York, NY: The Commonwealth Fund; 1999.

18. Mold J, Stein H. The cascade effect in the clinical care of patients. N Engl J Med. 1986;314:512-514.

19. Shi L, Starfield B. Income inequality, primary care, and health indicators. J Fam Pract. 1999;48:275-284.

Related Links:

» Medical Mistakes Kill 100,000 per year



Thank you, Dr. Mercola, for the courage to take on the good fight.

Doctors Are The Third Leading Cause of Death in the US, Killing 225,000 People Every Year

8 Points

Mark as Spam

Posted On Apr 08, 2008

This article is a good reason why one should avoid doctors and mainstream medical practice if at all possible. Prevention is the best medicine for taking care of your health and well-being. A lot of it is just simple common sense approaches concerning diet, exercise, sleep, stress reduction, sunlight, supplements, and avoiding environmental toxins. One must actively research, study and act apon these approaches for better health and not by taking a passive approach by simply going to the doctor a few times a year. You'll end up in the grave sooner than you think.

-🧲 Reply Mark as **0** Points Spam Posted On Jul 19, 2008

The troll needs to go to his nearest doctor so he can get some surgery and medication to make the credibility he

says it lack's! And then he can be one of the numbers!!!!!!!!!!



Gazer

rdoc

F Reply



Apprentice User, Joined On 3/2008 Send Message



Novice User, Joined On 10/2007

6

8

Send Message

Add as Friend

Add as Friend

	Posted On Nov 14, 2008	Wizard Bob
	If you add the number of deaths caused by the proper use of prescription drugs, then the American Health Care System becomes the leading cause of death in the US.	Send Message Add as Friend
Ī	 Mark as Reply Points Spam 	
Email this article to a friend 🗿 Share this article 🖪 t 📽		
Would you like to republish this article?		
	SECURE RECEVENDED RELEASED RELEASED RELEASED	
Privacy Policy Terms of Use Contact Us Testimonials 🔂 NEWSLETTER 🔂 BLOG		
© Copyright 2009 Dr. Joseph Mercola. All Rights Reserved. If you want to use this article on your site please click here. This content may be copied in full, with copyright, contact, creation and information intact, without specific permission, when used only in a not-for-profit format. If any other use is desired, permission in writing from Dr. Mercola is required.		

* These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure or prevent any disease. If you are pregnant, nursing, taking medication, or have a medical condition, consult your physician before using this product.

Doctors Are The Third Leading Cause of Death in the US, Killing 225,000 People Every Year